Mon. Not. R. Astron. Soc. **000**, 1–9 (2002) Printed 2 February 2008 (MN LaTeX style file v1.4)

arXiv:astro-ph/0208057v1 2 Aug 2002

Population effects on the red giant clump absolute magnitude: The K-band.

Maurizio Salaris[1] and Léo Girardi[2]
[1] *Astrophysics Research Institute, Liverpool John Moores University, Twelve Quays House, Egerton Wharf, Birkenhead CH41 1LD, UK*
[2] *Osservatorio Astronomico di Trieste, Via Tiepolo 11, I-34131 Trieste, Italy*

Accepted 2002 ???. Received 2002, June ??; in original form 2002 ???

ABSTRACT

We present a detailed analysis of the behaviour of the Red Clump K-band absolute magnitude ($M_K^{\rm RC}$) in simple and composite stellar populations, in light of its use as standard candle for distance determinations. The advantage of using $M_K^{\rm RC}$, following recent empirical calibrations of its value for the solar neighbourhood, arises from its very low sensitivity to the extinction by interstellar dust. We show that, as in the case of the V- and I-band results, $M_K^{\rm RC}$ is a complicated function of the stellar metallicity Z and age t. In general, $M_K^{\rm RC}$ is more sensitive to t and Z than $M_I^{\rm RC}$, for high t and low Z. Morever, for ages above $\sim$1.5 Gyr, $M_K^{\rm RC}$ decreases with increasing Z, the opposite behaviour with respect to $M_V^{\rm RC}$ and $M_I^{\rm RC}$.

We provide data and equations which allow the determination of the K-band population correction $\Delta M_K^{\rm RC}$ (difference between the Red Clump brightness in the solar neighbourhood and in the population under scrutiny) for any generic stellar population. These data complement the results presented in Girardi & Salaris (2001) for the V- and I-band. We show how data from galactic open clusters consistently support our predicted $\Delta M_V^{\rm RC}$, $\Delta M_I^{\rm RC}$ and $\Delta M_K^{\rm RC}$ values.

Multiband VIK population corrections for various galaxy systems are provided. They can be used in conjunction with the method devised by Alves et al. (2002), in order to derive simultaneously reddening and distance from the use of VIK observations of Red Clump stars. We have positively tested this technique on the Galactic globular cluster 47 Tuc, for which both an empirical parallax-based main sequence fitting distance and reddening estimates exist. We have also studied the case of using only V and I photometry, recovering consistent results for both reddening and distance. Application of this method to an OGLE-II field, and the results by Alves et al. (2002), confirm a LMC distance modulus of about 18.50, in agreement with the HST extragalactic distance scale zero-point.

Key words: distance scale – galaxies: stellar content – galaxies: individual (Large magellanic Cloud) – stars: Hertzsprung-Russell (HR) diagram – stars: horizontal branch – solar neighbourhood

1 INTRODUCTION

During the last few years a large body of work has been focused on the use of helium burning red clump (RC) stars as distance indicator; this has been mainly due to the fact that *Hipparcos* parallaxes allows an accurate calibration of the average RC brightness in the solar neighbourhood (as a comparison, *Hipparcos* parallaxes of the closest Cepheid and RR Lyrae stars have much larger errors, and the calibration of their absolute magnitude has provided conflicting results, see e.g. Feast & Catchpole 1997; Luri et al. 1998), and that the RC is easily recognizable in the colour-magnitude-diagram (CMD) of intermediate-old stellar populations.

The determination of the absolute magnitude of the local RC in a given pass-band λ, $M_{\lambda,\rm local}^{\rm RC}$, and the apparent magnitude $m_\lambda^{\rm RC}$ of the RC in a given stellar population is not difficult, since in both the *Hipparcos* database of nearby stars, and in CMDs covering even a small fraction of a nearby galaxy, one finds several hundreds of clump stars, easily identifiable from their CMD location. As proposed by Stanek & Garnavich (1998), a non-linear least-square fit of the function

© 2002 RAS

$$N(m_\lambda) = a + bm_\lambda + cm_\lambda^2 + d\exp\left[-\frac{(m_\lambda^{\rm RC} - m_\lambda)^2}{2\sigma_{m_\lambda}^2}\right] \qquad (1)$$

to the histogram of stars in the clump region per magnitude bin provides the value of $m_\lambda^{\rm RC}$ and its associated standard error. By applying this procedure to the *Hipparcos* database of nearby stars, the RC absolute brightness in the I-band (Cousins) $M_I^{\rm RC}$ has been determined with accuracy of hundredths of magnitude (Paczyński & Stanek 1998; Stanek, Zaritsky & Harris 1998).

Once the mean apparent magnitude of the RC in a given photometric band, $m_\lambda^{\rm RC}$, is measured in a nearby galaxy, its absolute distance modulus $\mu_0 = (m - M)_0$ is easily derived by means of

$$\mu_0 = m_\lambda^{\rm RC} - M_{\lambda,{\rm local}}^{\rm RC} - A_{m_\lambda} + \Delta M_\lambda^{\rm RC}. \qquad (2)$$

In this equation, A_{m_λ} is the interstellar extinction to the RC population of an external galaxy, and $\Delta M_\lambda^{\rm RC} = M_{\lambda,{\rm local}}^{\rm RC} - M_{\lambda,{\rm galaxy}}^{\rm RC}$ is the population effect, i.e. the difference of the mean RC absolute magnitude between the local and external samples of stars.

After early claims (Paczyński & Stanek 1998; Udalski et al. 1998; Stanek et al. 1998) based mainly on the constancy of $M_I^{\rm RC}$ with respect to the colour $(V-I)$, that the population correction $(\Delta M_I^{\rm RC})$ is negligible in the I-band, Cole (1998), Girardi et al. (1998, hereafter Paper I) and Girardi & Salaris (2001, hereafter Paper II) have conclusively demonstrated that this is not the case. The in-depth analysis we presented in Paper II has shown how theoretical stellar evolution models are able to reproduce the morphology and properties of the I-band RC brightness in Galactic open clusters, of the local *Hipparcos* RC as well as the RC in a sample of external galaxies, when using current estimates of the Star-Formation-History (SFR) and Age-Metallicity-Relation (AMR) of their stellar populations. The models predict a non-negligible dependence of $M_I^{\rm RC}$ on both metallicity and age, so that the population correction $\Delta M_I^{\rm RC}$ must be taken into account. In particular, the dependence on age is complex and non-monotonic. This implies that accurate RC distances can be determined only if one can estimate SFR and AMR for the stellar population under scrutiny. If there are no determinations of these two key parameters, errors up to ∼0.3 mag in the derived distance modulus are to be accepted.

In the specific case of the Large Magellanic Cloud (LMC) distance, that sets the zero point of the extragalactic distance scale, we obtained a population correction in the I-band $\Delta M_I^{\rm RC}$=+0.20. Unfortunately, the evaluation of the RC distance to the LMC is plagued by what possibly are uncertainties in the estimate of the interstellar extinction. Romaniello et al. (2000) and Sakai, Zaritsky and Kennicutt (2000) obtain, respectively, $I_0^{\rm RC} = 18.12 \pm 0.02$ and $I_0^{\rm RC} = 18.06 \pm 0.02$ from *HST* data, after deriving the extinction to individual stars from multiband colour fitting using theoretical model atmospheres results; these values, used in conjunction with the appropriate evolutionary correction and $M_I^{\rm RC} = -0.23\pm0.03$ for the local RC, provide $\mu_0 = 18.55\pm0.04$ and $\mu_0 = 18.49\pm0.04$, respectively. On the other hand Udalski (2000) has obtained $I_0^{\rm RC} = 17.94 \pm 0.05$, using different photometric data and determining the average stellar reddenings from the *COBE*/*DIRBE* reddening maps (Schlegel, Finkbeiner & Davis 1998), which gives a much shorter value, $\mu_0 = 18.37 \pm 0.06$.

To overcome uncertainties related to the extinction correction, Alves (2000) and Grocholski & Sarajedini (2002) have recently discussed the use of the RC in the K-band; the main advantage of working in this wavelength range is a very much reduced sensitivity to the extinction ($A_K \sim 0.2\ A_I$). Alves (2000) has derived $M_K^{\rm RC} = -1.61\pm0.03$ for the local RC stars with *Hipparcos* parallaxes and spectroscopic metallicities, and obtained a distance to the Galactic centre ($\mu_0 = 14.58\pm0.11$, or 8.24±0.42 Kpc) using the observed RC K-band magnitude for a sample of Baade's window RC stars with the same mean metallicity of the local sample; the underlying assumption is that there is no potential age-effect on the K-band RC level. Grocholski & Sarajedini (2002) have tested the behaviour of $M_K^{\rm RC}$ on a sample of single-age, single-metallicity stellar populations, constituted by 14 Galactic open clusters and 2 globular clusters with independent estimates of reddening and distances, finding a good agreement with the behaviour predicted by the theoretical models by Girardi et al. (2000), which are the same used in Paper I and Paper II. Moreover, Alves et al. (2002) have used simultaneously V, I and K photometry of LMC RC stars to determine both the LMC distance and the extinction for the observed RC population. They have demonstrated the need to apply population corrections to the RC absolute brightness determined from the local clump, otherwise a negative extinction is obtained; their final result $\mu_0 = 18.506 \pm 0.033$ for the LMC ($\mu_0 = 18.493 \pm 0.033 \pm 0.03$ random plus systematic error, after correcting for the location of the observed fields with respect to the LMC center, and including an estimate of the probable systematic errors involved) and $E(B-V) = 0.089\pm0.015$ was obtained making use of evolutionary corrections derived with the techniques and models presented in paper II.

In light of these results we consider important to investigate theoretically the properties of the RC in the K-band, as well as to assess the simultaneous reliability of the population corrections to the V-, I- and K-band for a given stellar population.

In Sect. 2 we discuss in detail the properties of the RC in the K-band and its dependence on age and metallicity, while in Sect. 3 we perform further tests to assess the reliability of the population corrections predicted by theory.

In Sect. 4 we present multiband population corrections for the galaxy systems discussed in Paper II; in addition, we will discuss the technique used by Alves et al. (2002) for their LMC distance determination, and apply it to the case of the Galactic globular cluster 47 Tuc (for which reddening estimates and an accurate empirical main-sequence fitting distance do exist) and to a field of the LMC with independent reddening determinations.

Our final conclusions are presented in Sect. 5.

2 THE THEORETICAL RED CLUMP LOCATION IN THE K-BAND

As in Paper I and Paper II we will base the discussion of the model behaviour on the Girardi et al. (2000) set of evolutionary tracks and isochrones. We remark that different models in the literature present systematic luminosity differences for the core helium burning (CHeB) stars that have passed through the helium flash – which are the stars belonging to the observed RC populations – due mainly to the different values of the helium core mass at the flash (see, e.g., Castellani et al. 2000; Salaris, Cassisi & Weiss 2002); however, the variation of their brightness with respect to age and metallicity is much more consistently predicted by theory (Castellani et al. 2000; Salaris et al. 2002). As in Paper II, the main results of this paper will be based on the strictly differential use of the model predictions.

We start our analysis by showing in Figure 1 the location of RC stars at the beginning of the He-burning phase on a magnitude-effective temperature plane, in simple stellar populations (single-age, single-metallicity) of various ages between 1 and 14 Gyr, and two different metallicities* ($Z = 0.001$ and a solar $Z = 0.019$); the behaviour of the bolometric luminosity as well as the V-, I- and K-band brightness is displayed. One striking feature is readily apparent; while the bolometric, V and I brightness at fixed age increase for decreasing metallicity, the K brightness is either approximately constant or even decreases with decreasing metallicity. The V and I magnitude follow (even if with different sensitivities) the behaviour of the bolometric magnitude, which is essentially determined by the variation with metallicity of the core mass at the He-ignition; the higher the metallicity, the lower the core mass, the lower the luminosity during the RC phase. However, in case of the K-band the situation is completely changed, due to the behaviour of BC_K.

Figure 2 displays the trend of BC_K with respect to metallicity and effective temperature, for the RC stars. It turns out that BC_K is almost unaffected by Z, but it is strongly sensitive to the value of $T_{\rm eff}$. Lower effective temperatures correspond to higher values of BC_K; since $M_K = M_{\rm bol} - BC_K$, cooler RC stars tend to be brighter. This sensitivity of BC_K to $T_{\rm eff}$ is strong enough to reverse the dependence of the stellar brightness on the He-core mass. As for the dependence of the K brightness on age, one can deduce from Figure 1 a large effect for high ages and a reduced one for lower ages, especially at high metallicities. This is a straightforward consequence of the $T_{\rm eff}$ location of the models as a function of their age; in fact, for increasing metallicity and decreasing age the scaling with respect to $T_{\rm eff}$ is reduced, thus producing a smaller variation of BC_K.

It is important to remark that the behaviour of BC_K shown in Figure 2 is not peculiar of the particular set of bolometric corrections adopted in our models (based on Kurucz 1993 models), but it is also found, e.g., in the empirical results by Montegriffo et al. (1998).

In order to explain the RC behaviour in composite stellar populations, like the solar neighbourhood and the LMC, we have to consider the mean RC brightness in simple stellar populations. It is obvious that the expected differential properties of the mean RC brightness will somehow mirror the results shown in Figure 1.



Figure 1. The location of RC stars at the beginning of the He-burning phase on an absolute magnitude–effective temperature plane, in simple stellar populations (single-age, single-metallicity) of various ages (ages of 2, 3, 4, 5, 6, 8, 9, 10, 11, 13, 14 Gyr moving clockwise; for each metallicity the oldest RC is also the faintest one) and two different metallicities ($Z = 0.001$ and $Z = 0.019$). The age sequences are shown for, from top to bottom, K, I, bolometric, and V magnitudes.

In Paper II we have discussed in detail how to derive the mean RC level for a given stellar population. The most detailed approach comes from a population synthesis algorithm; a synthetic CMD is produced for a given stellar population model, and then $M_\lambda^{\rm RC}$ is derived by fitting eq. 1 to the synthetic RC data. This is the method we have followed to produce the population corrections presented in Paper II and the ones we discuss also in this paper. However, it is very instructive to summarize a simpler approach presented in Paper II, since it shows clearly how to decompose a complex stellar population into its elementary 'simple' constituents.

For a given isochrone of age and metallicity (t, Z), one can perform the following integral over the isochrone section corresponding only to CHeB stars:

$$\langle M_\lambda(t,Z)\rangle = -2.5\log\left[\frac{1}{N_{\rm cl}(t,Z)}\int^{\rm CHeB}\phi(m_{\rm i})\,10^{-0.4M_\lambda}{\rm d}m_{\rm i}\right] \quad (3)$$

where M_λ is the absolute magnitude in the pass-band λ, $m_{\rm i}$ is the initial mass of the star at each isochrone point, and $\phi(m_{\rm i})$ is the Salpeter IMF (number of stars with initial mass in the interval $[m_{\rm i}, m_{\rm i} + {\rm d}m_{\rm i}]$). $N_{\rm cl}$ is the number of clump stars (at age t) per unit mass of stars initially born. It is simply given by the integral of the IMF by number, along the CHeB isochrone section, i.e.

$$N_{\rm cl}(t,Z) = \int^{\rm CHeB}\phi(m_{\rm i})\,{\rm d}m_{\rm i}\ . \quad (4)$$

From eq. 3, accurate values for the mean clump magnitudes can be determined, in case of a simple stellar population. These mean magnitudes enter straightforwardly in the computation of the mean clump magnitude for a given

---

* In the course of the paper we will also denote the metallicity of a given stellar population with the spectroscopic notation $[{\rm M/H}] = \log({\rm M/H})_{\rm star} - \log({\rm M/H})_\odot$, where M is the total metal abundance. [M/H] is related to Z by means of the approximate relation $[{\rm M/H}] = \log(Z/Z_\odot)$, where we assume $Z_\odot = 0.019$. In case of a scaled-solar metal distribution [M/H] = [Fe/H].

Table 1. $N_{\rm cl}$ and $\langle M_K \rangle$, as a function of age and metallicity, from Girardi et al. (2000) isochrones.

	$Z = 0.0004$		$Z = 0.001$		$Z = 0.004$		$Z = 0.008$		$Z = 0.019$		$Z = 0.03$	
t (Gyr)	$N_{\rm cl}$ (10^{-4})	$\langle M_K \rangle$	$N_{\rm cl}$ (10^{-4})	$\langle M_K \rangle$	$N_{\rm cl}$ (10^{-4})	$\langle M_K \rangle$	$N_{\rm cl}$ (10^{-4})	$\langle M_K \rangle$	$N_{\rm cl}$ (10^{-4})	$\langle M_K \rangle$	$N_{\rm cl}$ (10^{-4})	$\langle M_K \rangle$
0.5	26.30	−1.969	25.70	−2.175	31.60	−2.191	34.30	−2.047	30.60	−2.024	29.40	−2.118
0.6	29.40	−1.775	30.60	−2.072	37.00	−1.909	40.10	−1.770	36.10	−1.785	33.90	−1.900
0.7	32.30	−1.686	33.70	−1.941	41.40	−1.693	44.90	−1.599	40.20	−1.607	37.30	−1.733
0.8	36.80	−1.626	37.60	−1.802	44.30	−1.536	48.50	−1.460	43.90	−1.485	40.20	−1.610
0.9	38.00	−1.601	40.90	−1.665	47.20	−1.417	51.80	−1.350	46.50	−1.388	42.90	−1.505
1.0	43.50	−1.540	44.40	−1.543	50.10	−1.309	54.80	−1.270	48.30	−1.312	45.40	−1.422
1.1	44.30	−1.546	47.00	−1.456	52.80	−1.222	56.70	−1.209	49.90	−1.254	46.60	−1.364
1.2	41.50	−1.603	48.00	−1.415	53.10	−1.183	54.30	−1.247	49.40	−1.225	47.20	−1.328
1.3	31.00	−1.711	44.00	−1.478	52.20	−1.236	53.50	−1.363	48.40	−1.251	47.10	−1.333
1.4	20.70	−1.711	28.00	−1.634	51.60	−1.382	27.80	−1.516	48.20	−1.338	46.40	−1.370
1.5	19.40	−1.699	23.00	−1.619	29.60	−1.517	23.20	−1.559	28.20	−1.446	45.80	−1.496
1.6	18.70	−1.685	21.40	−1.620	22.50	−1.541	20.60	−1.576	25.10	−1.519	31.60	−1.590
1.7	17.80	−1.675	20.10	−1.616	19.70	−1.563	18.80	−1.596	22.20	−1.554	24.30	−1.576
1.8	16.90	−1.664	19.00	−1.612	18.20	−1.585	17.10	−1.606	19.50	−1.568	20.90	−1.588
1.9	16.00	−1.650	18.00	−1.607	16.60	−1.595	16.30	−1.617	17.40	−1.580	18.90	−1.615
2.0	15.50	−1.629	16.90	−1.595	16.20	−1.602	15.30	−1.627	16.80	−1.599	17.40	−1.638
2.2	14.80	−1.593	16.20	−1.572	15.50	−1.612	14.10	−1.624	15.50	−1.640	16.00	−1.675
2.4	13.50	−1.559	14.90	−1.550	14.80	−1.595	13.30	−1.619	14.70	−1.668	14.40	−1.682
2.6	13.00	−1.530	14.20	−1.528	14.10	−1.578	12.80	−1.606	13.50	−1.673	12.90	−1.681
2.8	12.50	−1.503	13.60	−1.508	13.30	−1.570	12.30	−1.591	12.30	−1.646	12.00	−1.685
3.0	11.50	−1.484	12.20	−1.491	12.90	−1.564	11.00	−1.590	11.00	−1.621	11.50	−1.690
3.2	11.10	−1.468	11.90	−1.478	12.50	−1.558	10.70	−1.588	10.80	−1.623	11.00	−1.700
3.4	10.70	−1.454	11.50	−1.466	11.80	−1.553	10.50	−1.585	10.70	−1.623	10.80	−1.698
3.6	10.30	−1.440	11.10	−1.455	11.50	−1.542	10.30	−1.584	10.20	−1.626	10.60	−1.696
3.8	9.65	−1.425	10.50	−1.444	11.10	−1.532	10.00	−1.577	9.54	−1.615	9.99	−1.699
4.0	9.38	−1.408	10.30	−1.432	10.80	−1.523	9.67	−1.567	9.09	−1.603	9.63	−1.693
4.3	8.98	−1.385	9.93	−1.414	9.55	−1.509	9.14	−1.553	8.42	−1.586	9.23	−1.688
4.6	8.58	−1.364	9.57	−1.398	9.21	−1.498	8.29	−1.541	7.88	−1.581	8.83	−1.683
4.9	8.04	−1.345	9.17	−1.384	8.87	−1.488	7.93	−1.535	7.72	−1.581	8.07	−1.675
5.2	7.80	−1.326	8.74	−1.368	8.54	−1.479	7.68	−1.527	7.56	−1.578	7.69	−1.664
5.5	7.55	−1.308	8.44	−1.352	7.72	−1.469	7.43	−1.520	7.39	−1.574	7.31	−1.653
6.0	7.15	−1.280	7.94	−1.328	7.36	−1.449	6.47	−1.508	6.38	−1.564	6.46	−1.637
6.5	6.75	−1.255	7.44	−1.306	7.01	−1.430	6.16	−1.488	6.04	−1.550	6.06	−1.628
7.0	6.46	−1.225	6.63	−1.282	6.65	−1.413	5.87	−1.472	5.71	−1.536	5.82	−1.621
7.5	6.22	−1.196	6.36	−1.257	6.23	−1.395	5.59	−1.457	5.32	−1.524	5.59	−1.614
8.0	5.97	−1.168	6.08	−1.233	5.99	−1.375	5.32	−1.446	5.12	−1.514	5.31	−1.611
9.0	5.49	−1.118	5.55	−1.190	5.52	−1.339	5.02	−1.414	4.77	−1.497	4.98	−1.590
10.0	4.85	−1.052	5.06	−1.137	5.14	−1.306	4.71	−1.385	4.48	−1.480	4.66	−1.571
11.0	4.71	−0.986	4.80	−1.084	4.91	−1.265	4.65	−1.354	4.31	−1.456	4.53	−1.577
12.0	4.56	−0.929	4.55	−1.037	4.69	−1.225	4.42	−1.309	4.09	−1.431	4.42	−1.572

galaxy model, of total age T. In fact, in this case one needs to perform the following integral:

$$\langle M_\lambda({\rm gal}) \rangle = \frac{1}{N_{\rm cl}({\rm gal})} \int_{t=0}^{T} N_{\rm cl}(t,Z)\, \psi(t)\, \langle M_\lambda(t,Z) \rangle \, {\rm d}t \ , \quad (5)$$

where

$$N_{\rm cl}({\rm gal}) = \int_{t=0}^{T} N_{\rm cl}(t,Z)\, \psi(t)\, {\rm d}t \ . \quad (6)$$

The function $\psi(t)$ is the SFR (in $M_\odot$ by unit time) at a moment t in the past, for the galaxy model considered; in addition, also the AMR $Z(t)$ should be specified. One can average the magnitudes in eq. 5, instead of luminosities as in the previous eq. 3; the reason is, as explained in Paper II, that in this way one obtains a quantity similar to the $M_\lambda^{\rm RC}$ derived by means of eq. 1. As discussed in Paper II, the population corrections evaluated with this approach are in very good agreement with the results obtained from the population synthesis approach; they are directly evaluated from the average RC brightness of the elementary simple populations, whose properties are highlighted in the following.

Table 1 presents the K-band mean absolute magnitude of clump stars, $\langle M_K \rangle$, as a function of age and metallicity, as derived from Girardi et al. (2000) isochrones by means of eq. 3, together with the values of $N_{\rm cl}(t,Z)$. This table can be used to easily derive the population correction in the K-band for any galaxy model. The same information is provided for the V- and I-band in Table 1 of Paper II. It is important to remark the pronounced maximum of $N_{\rm cl}$ for ages between 1 and 2 Gyr, which causes these relatively



Figure 2. Bolometric corrections for giants in the $K-$band, for both $Z = 0.019$ (dashed line), and $Z = 0.0004$ (solid line).



Figure 3. The behaviour of the clump as a function of age, for several metallicities between [M/H] $= -1.67$ and 0.20, (Z between 0.0004 and 0.03).

younger clump stars to be very numerous in galaxies with recent star formation. This aspect is thoroughly discussed in Paper II, and in Girardi (1999).

Figure 3 portrays the behaviour of the mean RC magnitude in simple stellar populations of different metallicities and ages, for the V, I and K passbands. In line with our desire to use the models only in a differential way, we have actually plotted the theoretical population correction (not the actual mean RC magnitude) for any given simple population considered. The theoretical counterpart of the local RC has been computed using the SFR and AMR by Rocha-Pinto et al. (2000a,b – see Paper II for further discussions on this issue).

It is evident that the behaviour of the population corrections – and therefore of the mean RC brightness – is strongly dependent on the photometric band used. For ages larger than about 1.5 Gyr $\Delta M_K^{\rm RC}$ increases for increasing metallicity (hence $M_K^{\rm RC}$ decreases), which is exactly the contrary of what happens to $\Delta M_V^{\rm RC}$ and $\Delta M_I^{\rm RC}$. If we consider the entire [M/H] range spanned by our models (from -1.7 up to 0.20), for ages below ∼4 Gyr $M_K^{\rm RC}$ is on average less sensitive to [M/H] than both $M_V^{\rm RC}$ and $M_I^{\rm RC}$. However, when the age is larger than this limit $M_K^{\rm RC}$ is more affected by [M/H] than $M_I^{\rm RC}$; for ages larger than ∼11 Gyr, $M_K^{\rm RC}$ is more metallicity-dependent than also $M_V^{\rm RC}$, at least in case of [M/H] lower than ~ -0.7. In the [M/H] interval between 0.0 and ~ -0.4 (this lower limit corresponds approximately to the average metallicity of the LMC clump stars) $M_K^{\rm RC}$ is less affected by the metallicity than both $M_I^{\rm RC}$ and $M_V^{\rm RC}$, at least up to ages of about 11 Gyr. At super-solar [M/H] and for ages above ∼3 Gyr, $M_K^{\rm RC}$ is more sensitive to metallicity than the V and I brightness.

This complex relationship among the RC V, I and K brightness stems from the different sensitivities to age. In particular, $M_K^{\rm RC}$ is more affected by age than both $M_I^{\rm RC}$ and $M_V^{\rm RC}$, for the lower metallicities and higher ages.

3 TESTING THE THEORETICAL MULTICOLOUR POPULATION CORRECTIONS

Paper II and Grocholski & Sarajedini (2002) have already shown separately the consistency of the theoretical results in the I and K band, when applied to a sample of Galactic open clusters of various ages and metallicities. The very different behaviour of the population corrections in different passbands allows one to perform the following additional test for the accuracy of the models.

Figure 4 compares the difference of the population corrections in two different photometric bands, between theory and observations, for a sample of Galactic open clusters with photometry in V, I and K. The data about the clusters V and I magnitudes, as well as ages, [Fe/H] and reddening values are taken from Sarajedini (1999). K-band photometric data are from Grocholski & Sarajedini (2002). The clusters' ages have been derived with respect to the reference cluster M 67, assumed to be 4 Gyr old on the base of previous studies; the age scaling has been obtained from the distance modulus difference (derived from the Main Sequence-fitting) with respect to M 67. It is important to notice that the difference of the evolutionary corrections in two different photometric bands is independent of the adopted distance modulus (while the cluster age is dependent on the relative Main Sequence-fitting distance moduli and the age zero-point). As for the empirical calibration of the local RC we have used the values provided by Alves et al. (2002), namely $M_K^{\rm RC} = -1.60 \pm 0.03$, $M_I^{\rm RC} = -0.26 \pm 0.03$, $M_V^{\rm RC} = 0.73 \pm 0.03$ (the theoretical values we obtain from our simulation are $M_K^{\rm RC} = -1.54$, $M_I^{\rm RC} = -0.17$, $M_V^{\rm RC} = 0.84$; we stress again that nowhere we use individual absolute magnitudes



Figure 4. The population correction difference between two different photometric bands, as predicted by theory (lines) and from observations of a sample of galactic open clusters (dots) with photometry in V, I and K. The theoretical metallicities are [M/H] = −0.38 (long-dashed line), 0 (dot-short-dashed), and 0.20 (dot-long-dashed). The numbers on the plot indicate the metallicity [M/H] for each cluster; the clusters are, from left to right, NGC 2204, NGC 6819, M 67, Be 39 and NGC 6791.

from the stellar models, but only magnitude differences). The value $M_I^{\rm RC} = -0.26 \pm 0.03$ agrees, within 1σ, with the previous determination by Stanek & Garnavich (1998).

The agreement between theory and observations is satisfactory – also considering the uncertainty of the order of ∼0.1 dex associated to the individual cluster [M/H] estimates –, another confirmation for the population corrections predicted by the stellar models we employ. In fact, taking into account the very different behaviour of the predicted evolutionary corrections in V, I and K, it is highly improbable that a combination of errors in the theory and in the adopted clusters' parameters, were to produce such a good agreement.

Another interesting test makes use of the available V- and K-photometry of the few RC stars populating the *Hipparcos* database for the Hyades and Praesepe clusters. There is a large body of work suggesting that these two clusters share the same age and metallicity (e.g., the discussion and references in van Leeuwen 1999; Castellani et al. 2002). From the *Hipparcos* data the cluster distances result to be $\mu_0^{\rm Hyades} = 3.33 \pm 0.01$ and $\mu_0^{\rm Praesepe} = 6.37 \pm 0.15$ (Perryman et al. 1997; van Leeuwen 1999); a comparison of their CMD corrected for the different distance moduli, as performed by Castellani et al. (2002), confirms a basically identical metallicity and age, since the good overlap of the upper main sequence and turn off, and also of the RC region.

For a common age $t = 625 \pm 50$ million years (higher than the lower limit for the existence of the RC, which is of about 500 million years) and [Fe/H] = 0.14 (Perryman et al. 1997), we have determined the appropriate population correction and plotted in Figure 5 the theoretical luminosity function (number of stars per magnitude bin) for RC stars in the V- and K-band (the normalization is arbitrary); the brightness of the RC has the appropriate mean value obtained after applying the theoretical population corrections (which are of the order of ∼ −0.3 mag in both passbands) to the observed local RC absolute magnitude. On the same plot we show the V and K absolute magnitude of the 6 stars (2 in the Hyades and 4 in Praesepe) populating the RC of the composite CMD of these two clusters. Due to the small size of the star sample it is not possible to determine a meaningful mean brightness to compare with theory, but one can notice that the empirical stellar absolute magnitudes are in good agreement with the location and the width of the RC as predicted by the theoretical population corrections.



Figure 5. The theoretical luminosity function (number of stars per magnitude bin) for Hyades RC stars in the V- and K-band (the normalization is arbitrary); the brightness of the RC has the appropriate mean value obtained after applying the theoretical population corrections (which are of the order of ∼ −0.3 mag in both passbands) to the local RC absolute magnitude. On the same plot we show the V and K absolute magnitude of the 6 stars (2 in the Hyades and 4 in Praesepe; open and closed circles, respectively) populating the RC of the composite CMD of these two clusters (see text for details).

As a warning, it is also important to take into account the fact that the values of $\Delta M_K^{\rm RC}$ and $\Delta M_V^{\rm RC}$ are both strong function of t in this age range (in case Praesepe is younger than the Hyades by ∼ 100 million years, its RC would be on average brighter by ∼0.2 mag than the Hyades RC); we notice however that here is no obvious offset between the two sets of observational points, and between them and the theoretical luminosity function. In conclusion, we regard the agreement between theoretical and empirical data in Figure 5 as an additional indication of the accuracy of the population corrections predicted by theory.

The generally small values of $\Delta M_K^{\rm RC}$ for ages between ∼1.5 and ∼4 Gyr, as displayed in Figure 3, raise the question of to what extent one is allowed to disregard the population corrections when considering the K-band RC magnitude as



Figure 6. The population corrections to the K-band for simple stellar populations of varying age and metallicity, marking the region for which the population correction is within ±0.1 mag.



Figure 7. Empirical CMDs for the local RC (from Alves 2000) in the V, I and K passbands.

distance indicator (see also the discussion in Grocholski & Sarajedini 2002). In this age range, the lower sensitivity of the K-band RC brightness to both age and metallicity may suggest to use the local K-band RC brightness as a perfect standard candle, without taking into account the appropriate $\Delta M_K^{\rm RC}$.

To answer this question, Figure 6 shows again the population corrections to the K band for simple stellar populations, but this time it is also marked the region in which the population correction is within ±0.1 mag. If a systematic uncertainty of this order on the RC distance is considered to be acceptable, the allowed range of age and metallicity is easily evaluated. For example, one can obtain distances with this precision, without evaluating the population correction, for t and Z values typical of the sample of Galactic open clusters older than ∼2.5 Gyr. It is however not allowed to neglect *apriori* the evaluation of $\Delta M_K^{\rm RC}$, due to the fact that it can reach values as high as ±0.5 mag.

4 *V*-,*I*-, *K*-BAND POPULATION CORRECTIONS FOR EXTERNAL GALAXIES

Having assessed the adequacy of the theoretical population corrections for simple stellar populations, in this section we proceed to compute the corrections in V, I and K for the composite systems discussed in Paper II, using the appropriate SFR and AMR discussed in Paper II; the results are summarized in Table 2. The full references for the SFR and AMR can be found in table 4 of Paper II.

As a test, we show in Figs. 7 and 8 the empirical data for the local RC (from Alves 2000) in the V, I and K passbands, and the results from the theoretical simulations, respectively. Notice the good correspondence between the predicted and observed shape of the RC in the different colour planes. In particular, the RC in the M_I–$(V-I)$ plane is basically horizontal (see the in-depth discussion of Paper II on the local RC and its properties in the M_I–$(V-I)$ plane), while in the M_K–$(V-K)$ plane is slightly tilted towards higher M_K values, and in the M_V–$(V-I)$ plane is tilted towards lower M_V values. These properties are well reproduced by the theoretical models. More in detail, the observed slopes $\Delta M_V/\Delta(V-I)$ and $\Delta M_K/\Delta(V-K)$ are equal to, respectively, 1.46±0.27 and −0.63±0.12, which compare well with the corresponding theoretical values 1.18±0.07 and −0.54±0.02. In addition, the theoretical simulation predicts no significant correlations between [Fe/H] and either M_K or M_V, in agreement with the empirical data by Alves (2000).

The data in Table 2, in conjunction with the empirical V, I and K absolute brightness of the local RC reported in Section 3, can be used to derive simultaneously reddening and distances to nearby galaxies, following the technique employed by Alves et al. (2002). The basic idea is that the apparent distance moduli determined simultaneously in various photometric bands, using the appropriate population corrections, must all provide the same unreddened distance; after assuming a reddening law, one can therefore use this constraint to derive simultaneously reddening and true distance modulus. Alves et al. (2002) have adopted for the reddening law the following ratios: A_K/E(B-V)=0.35, A_I/E(B-V)=1.96, A_V/E(B-V)=3.24. We will use these same ratios throughout the rest of this section.

The LMC value of $\Delta M_V^{\rm RC}$ provided in Table 2 is slightly different from the one used by Alves et al. (2002), which we obtained after a preliminary evaluation of this quantity. This small difference (0.04 mag) does not influence appreciably the results obtained by Alves et al. (2002). In fact, we repeated their analysis employing the $\Delta M_V^{\rm RC}$ of Table 2 ($\Delta M_I^{\rm RC}$ and $\Delta M_K^{\rm RC}$ are unchanged), and obtained μ_0=18.505±0.045 and $E(B-V)$=0.079±0.014, almost coin-



Figure 8. The same as Fig. 7, but for a theoretical simulation with a much larger number of stars.

Table 2. Population corrections for the clump in nearby galaxy systems.

System	$\Delta M_V^{\rm RC}$	$\Delta M_I^{\rm RC}$	$\Delta M_K^{\rm RC}$
Solar Neighbourhood (*Hipparcos*)	0.00	0.00	0.00
Baade's Window (scaled-solar)	−0.21	−0.08	−0.07
Baade's Window (α-enhanced)	−0.06	−0.01	−0.11
Carina dSph	+0.59	+0.35	−0.17
SMC	+0.31	+0.29	−0.07
LMC	+0.26	+0.20	−0.03

cident with μ_0=18.506±0.033 and $E(B-V)$=0.089±0.015 obtained by Alves et al. (2002).

We have further tested this method (and the corresponding population corrections) on the Galactic globular cluster 47 Tuc, for which an accurate *Hipparcos*-based empirical main-sequence fitting distance μ_0=13.25±0.07 has been recently provided by Percival et al. (2002), and reddening estimates are also available ($E(B-V)$=0.04±0.02, see the discussion in Percival et al. 2002). Spectroscopic analyses give [Fe/H]=−0.70±0.1, together with an enhancement of the α-elements typical of the Galactic Halo population. We have taken into account the α-element enhancement – as in Paper II – using the results by Salasnich et al. (2000). V- and I-band data come from the photometry employed by Percival et al. (2002), while K-band data are taken from the analysis by Grocholski & Sarajedini (2002). We have assumed an *a priori* age of 11 ± 2 Gyr for the cluster, which corresponds to the age range allowed by an uncertainty of more than ±0.1 mag around the main sequence fitting distance. By applying Alves et al. (2002) procedure we have obtained μ_0=13.19±0.07 and $E(B-V)$=0.055±0.015, in good agreement with the independent determinations given before.

K-band data for a large variety of Galactic and extragalactic stellar populations are not yet widely available, while VI data are much more common in the literature. We have therefore applied Alves et al. (2002) technique using only VI data. If, as shown in the previous section, our population corrections are accurate, one expects to obtain distances consistent with the results from VIK data, albeit possibly with larger errors due to the lack of an additional constraint from the K-band magnitudes. In case of old metal poor stellar populations, the RC V and I brightness is much less dependent on age than the K magnitude, so that the assumptions about the stellar ages are less critical. By repeating the determination of 47 Tuc distance with the VI data only, we obtained μ_0=13.26±0.07 and $E(B-V)$=0.02±0.02, consistent with the result from VIK photometry. The same exercise can be performed on Alves et al. (2002) LMC data; we derived in this case μ_0=18.53±0.07 and $E(B-V)$=0.071±0.022, in very good agreement with the result from the full VIK analysis.

As a final exercise, we have studied the field LMC-SC6 from the OGLE-II database (Udalski et al. 2000); this field overlaps with the field centred on the eclipsing binary HV982, studied by Larsen et al. (2000). We have corrected the zero point of the OGLE-II VI photometry, in order to place it on the same system of Alves et al. (2002), and employed stars with random photometric error smaller than 0.04 mag.

We obtain for the RC level in field LMC-SC6 $V^{\rm RC}$=19.21 and $I^{\rm RC}$=18.17. By employing the population corrections given in Table 2 we have obtained μ_0=18.47±0.04 and $E(B-V)$=0.085±0.015. This distance agrees well within the errors with Alves et al. (2002) results. The minimum reddening for this field as estimated by Larsen et al. (2000) is $E(B-V)$=0.085±0.02. Fitzpatrick et al. (2002) obtain from spectral fitting of the eclipsing binary HV982 a value $E(B-V)$=0.086±0.005. Oestreicher et al. (1995) provides a minimum reddening of 0.05 mag. All these values are consistent with our reddening estimate for the RC population.

5 CONCLUSIONS

We have presented a detailed analysis of the behaviour of the RC brightness in the K-band, together with tests for the accuracy of the predicted population corrections to the local RC VIK brightness, and a discussion of the Alves et al. (2002) method for deriving simultaneously distance and reddening using multiband RC photometric data. More in detail:

– we have shown that the K-band brightness of the RC is mainly determined by the behaviour of the bolometric correction to the K-band. For ages larger than about 1.5 Gyr the trend of the RC brightness with respect to [M/H] is reversed in comparison with the V- and I-band, that is, $M_K^{\rm RC}$ decreases for increasing metallicity. By considering the entire [M/H] range spanned by our models (Girardi et al. 2000), for ages below ∼4 Gyr $M_K^{\rm RC}$ is on average less sensitive to [M/H] than both $M_V^{\rm RC}$ and $M_I^{\rm RC}$. At higher ages $M_K^{\rm RC}$ is more affected by [M/H] than $M_I^{\rm RC}$; for ages larger than ∼11 Gyr it is more metallicity-dependent than also $M_V^{\rm RC}$, at least for [M/H] lower than ~ -0.7. It is interesting to notice that for [M/H] between 0.0 and ~ -0.4 $M_K^{\rm RC}$ is less affected by the metallicity than both $M_I^{\rm RC}$ and $M_V^{\rm RC}$, at least up to ages of about 11 Gyr. This complex relationship among the RC V, I and K brightness is due to their different sensitivities to age. In particular, $M_K^{\rm RC}$ is more affected by age than both $M_I^{\rm RC}$ and $M_V^{\rm RC}$, for the lower metallicities and higher ages. Due to the non-trivial dependence of $M_K^{\rm RC}$ on both [M/H] and age, it is always necessary – as in the case of $M_V^{\rm RC}$ and $M_I^{\rm RC}$ – to determine the appropriate population correction for the stellar population under scrutiny, when using the RC as a standard candle;

– we have provided in Table 1 values of the RC average K brightness for a large range of [M/H] and ages. These data, used in conjunction with eq. 5, eq. 6, and the data in Table. 1 of Paper II, can be employed to determine multiband population corrections for any given stellar population, once SFR and AMR are prescribed;

– we have simultaneously and positively tested the evolutionary corrections $\Delta M_V^{\rm RC}$, $\Delta M_I^{\rm RC}$ and $\Delta M_K^{\rm RC}$ against Galactic open cluster data;

– we have discussed the method applied by Alves et al. (2002) for determining reddening and distance from the V-, I- and K-band RC brightness; we have positively tested this technique on the Galactic globular cluster 47 Tuc, for which both an empirical parallax-based main sequence fitting distance and reddening estimates exist. We have also studied the case of using only V and I photometry, recovering consistent results for both reddening and distance. Application of this method to an OGLE-II field, and the results by Alves et al. (2002), confirm a LMC distance modulus of about 18.50, in agreement with the *HST* extragalactic distance scale zero-point (Freedman et al. 2001).

ACKNOWLEDGMENTS

We thank D. Alves for his useful comments on a preliminary version of the manuscript.

REFERENCES

Alves D., 2000, ApJ, 539, 732

Alves D., Rejkuba M., Minniti D., Cook K.H., 2002, ApJ 573, L51

Castellani V., Degl'Innocenti S., Girardi L., Marconi M., Prada Moroni P.G., Weiss A., 2000, A&A 354, 350

Castellani V., Degl'Innocenti S., Prada Moroni P.G., Tordiglione V., 2002, MNRAS 334, 193

Cole A.A., 1998, ApJ 500, L137

Feast M.W., Catchpole R.M., 1997, MNRAS 286, L1

Fitzpatrick E.L., Ribas I., Guinan E.F., DeWarf L.E., Maloney F.P., Massa D., 2002, ApJ 564, 260

Freedman W., et al., 2001, ApJ 533, 47

Girardi L., 1999, MNRAS 308, 818

Girardi L., Groenewegen M.A.T., Weiss A., Salaris M., 1998, MNRAS 301, 149 (Paper I)

Girardi L., Bressan A., Bertelli G., Chiosi C., 2000, A&AS, 141, 371

Girardi L., Salaris M., 2001, MNRAS 323, 109 (Paper II)

Grocholski A.J., Sarajedini A., 2002, AJ 123, 1603

Kurucz R.L., 1993, in IAU Symp. 149: The Stellar Populations of Galaxies, eds. B. Barbuy, A. Renzini, Dordrecht, Kluwer, p. 225

Larsen S.S., Clausen J.V., Storm J., 2000, A&A 364, 455

Luri X., Gomez A.E., Torra J., Figueras F., Mennessier, M.O., 1998, A&A 335, L81

Montegriffo P., Ferraro F.R., Origlia L., Fusi Pecci F., 1998, MNRAS 297, 872

Oestreicher M.O., Gochermann J., Schmidt-Kaler T., 1995, A&AS 112, 495

Paczyński B., Stanek K.Z., 1998, ApJ 494, L219

Percival S.M., Salaris M., van Wyk F., Kilkenny D., 2002, ApJ 573, 174

Perryman M.A.C., et al., 1997, A&A 323, L49

Rocha-Pinto H.J., Maciel W.J, Scalo J., Flynn C., 2000a, A&A 358, 850

Rocha-Pinto H.J., Scalo J., Maciel W.J, Flynn C., 2000b, A&A 358, 869

Romaniello M., Salaris M., Cassisi S., Panagia N. 2000, ApJ 530, 738

Sakai S., Zaritsky D., Kennicutt R.C. Jr., 2000, AJ 119, 1197

Salaris M., Cassisi S., Weiss A., 2002, PASP 114, 375

Salasnich B., Girardi L., Weiss A., Chiosi C., 2000, A&A 361, 1023

Sarajedini A., 1999, AJ 118, 2321

Schlegel D.J., Finkbeiner D.P., Davis M., 1998, ApJ 500, 525

Stanek K.Z., Garnavich P.M., 1998, ApJ 503, L131

Stanek K.Z., Zaritsky D., Harris J., 1998, ApJ 500, L141

Udalski A., Szymański M., Kubiak M., Pietrzyński G., Woźniak P., Żebruń K., 1998, Acta Astr. 48, 1

Udalski A., Szymański M., Kubiak M., Pietrzyński G., Soszynski I., Wozniak, P., Żebruń K., 2000, Acta Astr. 50, 307

Udalski A., 2000, ApJ 531, L25

van Leeuwen F., 1999, A&A 341, L71

Zaritsky D., 1999, AJ 118, 2824